


CHEMICAL COMPOSITION

Active ingredient: Diphacinone
(2-Diphenyl acetyl-1, 3-indandione) 0.005%
LD₅₀: 10mg/kg (Rat oral dose).

STRUCTURAL FORMULA



PRECAUTION

- Do not use the product in the place where children or domestic animals can eat it by mistake.
- Please use gloves for wild rodents avoid human odor.
- Be careful to use the amount prescribed at the back.
- It is effective to put the product on dumps, roots of trees or circumferences of the place where rodents are frequent.


ANTIDOTE

If swallowed the product, drink 1 or 2 glasses of water and induce vomiting by touching back of throat with a clean finger, then call a physician.

If the prothrombin time extends over as twice as its normal level (ca. 15sec.), oral dose of Vitamin K₁ is indicated.



FOR BETTER PROFIT IN YOUR FIELD YASODION®

NEW POWERFUL RODENTICIDE



OHTSUKA CHEMICAL INDUSTRIAL CO., LTD.

3-25 NISHI-IKEBUKURO, TOSHIMA-KU, TOKYO, JAPAN



CHARACTERISTICS

- ★ Excellent efficacy Good poisoning effect at lower dosage.
- ★ Very palatable No bait shyness. Mice and rats eat YASODION willingly.
- ★ Easy to use Ready for use anytime. No preparation is necessary.
- ★ Good for a long time use Waterproof bag keeps YASODION from rain and moisture for better and long performance.
- ★ Safety YASODION is safe to domestic and wild animals under usual use.※

DIFFERENCE BETWEEN YASODION AND OTHER RODENTICIDES

- 1) YASODION is an anticoagulant poison and quite effective at lower concentration than other rodenticides (0.005% of diphacinone).
- 2) YASODION contains attractive ingredients so mice and rats show no bait shyness to it.
- 3) YASODION is stable in rain because it is packed in waterproof bags, while others are not protected from rain and moisture.

Our special formulation makes YASODION the best in total performance.


※Lethal Dose of YASODION

Lethal Dose 50 (LD50)

ANIMAL weight	RAT 250g	CAT 2.5kg	DOG 5.0kg	PIG 100kg	BIRD 1.0kg
YASODION	16~25.5g	735g	300~750g	300kg	>40kg

DIRECTIONS FOR USE

- In paddy fields
Put 1 or 2 bags of YASODION in a burrow or on a pathway of rodents at the rate of 20~30 bags per 10a (1000m²).
- In sugarcane or corn field
Put 1 or 2 bags of YASODION every 10m in the field and the circumference.
The rate of 20~30 bags per 10a (1000m²).



● Bait point in field.
○ Bait point in circumference.

